

Title	On the non-homogeneous central Morrey type spaces in $L^1(\mathbf{R}^n)$ and the weak boundedness of some operators (The deepening of function spaces and its environment)
Author(s)	Matsuoka, Katsuo; Mizuta, Yoshihiro
Citation	数理解析研究所講究録 = RIMS Kokyuroku (2018), 2095: 88-96
Issue Date	2018-12
URL	http://hdl.handle.net/2433/251709
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

On the non-homogeneous central Morrey type spaces in $L^1(\mathbf{R}^n)$ and the weak boundedness of some operators

Katsuo MATSUOKA and Yoshihiro MIZUTA

Dedicated to the memory of Professor Yasuji Takahashi

Abstract

Our aim in this note is to discuss the weak boundedness of the maximal and generalized Riesz potential operators in the non-homogeneous central Morrey type space $M^{1,q,\nu}(\mathbf{R}^n)$ of all measurable functions f on \mathbf{R}^n satisfying

$$\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)} = \left(\int_1^\infty (r^{-\nu} \|f\|_{L^1(B(0,r))})^q \frac{dr}{r} \right)^{1/q} < \infty$$

for $-\infty < \nu < \infty$ and $0 < q \leq \infty$; when $q = \infty$, we apply a necessary modification. To do this, we consider the family $WM^{p,q,\nu}(\mathbf{R}^n)$ of all functions $f \in L^p_{loc}(\mathbf{R}^n)$ such that

$$\|f\|_{WM^{p,q,\nu}(\mathbf{R}^n)} = \sup_{\lambda > 0} \int_1^\infty \left(r^{-\nu} \lambda |\{x \in B(0,r) : |f(x)| > \lambda\}|^{1/p} \right)^q \frac{dr}{r} < \infty,$$

where $1 \leq p \leq \infty$.

1 Introduction

In the n -dimensional Euclidean space \mathbf{R}^n , the space $B^p(\mathbf{R}^n)$ given by Beurling [5] is a special case of Herz spaces $K_p^{\alpha,r}(\mathbf{R}^n)$ introduced by Herz [13]. As an extension of the space $B^p(\mathbf{R}^n)$, Alvarez, Guzmán-Partida and Lakey [4] introduced the non-homogeneous central Morrey space $B^{p,\nu}(\mathbf{R}^n)$. Fu, Lin and Lu [11] proved the boundedness of the Riesz potential operator I_α on $B^{p,\nu}(\mathbf{R}^n)$, where $-n/p \leq \nu < -\alpha$; see also [16].

It is well known that the maximal operator M is weakly bounded in the Lebesgue space $L^1(\mathbf{R}^n)$ (see [23]). We extend it to the non-homogeneous central Morrey type space $M^{1,q,\nu}(\mathbf{R}^n)$ consisting of all measurable functions f on \mathbf{R}^n satisfying

$$\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)} = \left(\int_1^\infty (r^{-\nu} \|f\|_{L^1(B(0,r))})^q \frac{dr}{r} \right)^{1/q} < \infty$$

2000 Mathematics Subject Classification : Primary 31B15, 46E35

Key words and phrases : non-homogeneous central Morrey type space, maximal function, grand Morrey spaces in L^1 , generalize Riesz potentials, Sobolev's inequality

The first author is partially supported by Grant-in-Aid for Scientific Research (C), No. 17K05306, Japan Society for the Promotion of Science and Grant 2017 Overseas Research of the College of Economics, Nihon University, Japan, and the second author is partially supported by Grant-in-Aid for Scientific Research (C), No. 26400154, Japan Society for the Promotion of Science.

when $0 < q < \infty$ and

$$\|f\|_{M^{1,\infty,\nu}(\mathbf{R}^n)} = \sup_{r>1} r^{-\nu} \|f\|_{L^1(B(0,r))} < \infty$$

when $q = \infty$, where $B(x, r)$ denotes the open ball centered at x of radius r .

There are several Morrey type spaces related to our non-homogeneous central Morrey type spaces; e.g., Morrey spaces by Adams-Xiao [1], local Morrey type spaces and complementary local Morrey type spaces by Burenkov and al. [3], [6, 7], [8], [9], grand Morrey spaces by Kokilashvili-Meskhi-Rafeiro [12] (see also [10], [17]), and Herz-Morrey spaces by the second author and Ohno [20, 21].

Our aim in this note is to establish the weak boundedness of the maximal and generalized Riesz potential operators in $M^{1,q,\nu}(\mathbf{R}^n)$.

2 Non-homogeneous central Morrey type spaces

DEFINITION 2.1 (Non-homogeneous central Morrey type spaces). For $1 \leq p \leq \infty$, $0 < q \leq \infty$ and $-\infty < \nu < \infty$, we define a non-homogeneous central Morrey type spaces $M^{p,q,\nu}(\mathbf{R}^n)$ of all measurable functions f on \mathbf{R}^n such that

$$\|f\|_{M^{p,q,\nu}(\mathbf{R}^n)} = \left(\int_1^\infty (r^{-\nu} \|f\|_{L^p(B(0,r))})^q \frac{dr}{r} \right)^{1/q} < \infty$$

when $q < \infty$ and

$$\|f\|_{M^{p,\infty,\nu}(\mathbf{R}^n)} = \sup_{r>1} r^{-\nu} \|f\|_{L^p(B(0,r))} < \infty$$

when $q = \infty$.

Note that for $1 \leq p \leq \infty$,

- (1) if $\nu = 0$, then $M^{p,\infty,\nu}(\mathbf{R}^n) = L^p(\mathbf{R}^n)$;
- (2) if $\nu < 0$, then $M^{p,\infty,\nu}(\mathbf{R}^n) = \{0\}$;
- (3) if $\nu > 0$, then $M^{p,\infty,\nu}(\mathbf{R}^n) \supset L^p(\mathbf{R}^n)$.

For fundamental properties of our Morrey type spaces, we have the following.

LEMMA 2.2. Let $1 \leq p \leq \infty$ and $-\infty < \nu < \infty$. For $0 < q_1 < q_2 < \infty$,

$$M^{p,q_1,\nu}(\mathbf{R}^n) \subset M^{p,q_2,\nu}(\mathbf{R}^n) \subset M^{p,\infty,\nu}(\mathbf{R}^n).$$

LEMMA 2.3. If $1 \leq p \leq \infty$, $-\infty < \nu < \infty$ and $0 < q < \infty$, then

$$\|f\|_{M^{p,q,\nu}(\mathbf{R}^n)} \sim \left(\sum_{j=1}^{\infty} (2^{-\nu j} \|f\|_{L^p(B(0,2^j))})^q \right)^{1/q},$$

where the symbol $g \sim h$ means that $C^{-1}h \leq g \leq Ch$ for some constant $C > 0$.

3 Maximal functions

For a locally integrable function f on \mathbf{R}^n , the maximal function of f is defined by

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x, r)|} \int_{B(x, r)} |f(y)| dy,$$

where $|B(x, r)|$ denotes the Lebesgue measure of the open ball $B(x, r)$ centered at $x \in \mathbf{R}^n$ of radius r . It is well known that the maximal operator $M : f \rightarrow Mf$ is weakly bounded in $L^1(\mathbf{R}^n)$, that is,

$$|\{x \in \mathbf{R}^n : Mf(x) > \lambda\}| \leq C\lambda^{-1} \int_{\mathbf{R}^n} |f(y)| dy$$

for all $\lambda > 0$ and $f \in L^1(\mathbf{R}^n)$.

DEFINITION 3.1 (Weak central Morrey type spaces). For $1 \leq p \leq \infty$, $0 < q \leq \infty$ and $-\infty < \nu < \infty$, we denote by $WM^{p,q,\nu}(\mathbf{R}^n)$ the family of all functions $f \in L^p_{loc}(\mathbf{R}^n)$ such that

$$\|f\|_{WM^{p,q,\nu}(\mathbf{R}^n)} = \sup_{\lambda>0} \left(\int_1^\infty (r^{-\nu} \lambda |\{x \in B(0, r) : |f(x)| > \lambda\}|^{1/p})^q \frac{dr}{r} \right)^{1/q} < \infty$$

when $q < \infty$ and

$$\|f\|_{WM^{1,\infty,\nu}(\mathbf{R}^n)} = \sup_{\lambda>0, r>1} r^{-\nu} \lambda |\{x \in B(0, r) : |f(x)| > \lambda\}|^{1/p} < \infty$$

when $q = \infty$.

In view of A. Almeida and D. Drihem [2], we know that the maximal operator M is bounded in $M^{p,q,\nu}(\mathbf{R}^n)$, when $1 < p < \infty$. The case $p = 1$ is treated in the following.

THEOREM 3.2. Let $0 \leq \nu < n$ and $0 < q \leq \infty$, or let $\nu = n$ and $q = \infty$. Then the maximal operator M is bounded from $M^{1,q,\nu}(\mathbf{R}^n)$ to $WM^{1,q,\nu}(\mathbf{R}^n)$, that is, there exists a constant $C > 0$ such that

$$\|Mf\|_{WM^{1,q,\nu}(\mathbf{R}^n)} \leq C \|f\|_{M^{1,q,\nu}(\mathbf{R}^n)}$$

for $f \in M^{1,q,\nu}(\mathbf{R}^n)$.

Proof. We show only the case when $1 < q < \infty$, because the remaining case is easily obtained.

Let f be a measurable function on \mathbf{R}^n such that $\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)} \leq 1$. For $r > 1$, we write

$$f = f\chi_{B(0,2r)} + f\chi_{\mathbf{R}^n \setminus B(0,2r)} = f_1 + f_2,$$

where χ_E denotes the characteristic function of a measurable set $E \subset \mathbf{R}^n$. Note here that

$$\begin{aligned} Mf_2(x) &\leq \sup_{t \geq r} \frac{1}{|B(0, t)|} \int_{B(0,2t) \setminus B(0,2r)} |f(y)| dy \\ &\leq C \int_{\mathbf{R}^n \setminus B(0,2r)} |f(y)| |y|^{-n} dy \end{aligned}$$

for $x \in B(0, R)$. Let $\lambda > 0$. Since

$$\begin{aligned} & \{x \in B(0, r) : Mf(x) > \lambda\} \\ & \subset \{x \in B(0, r) : Mf_1(x) > \lambda/2\} \cup \{x \in B(0, r) : Mf_2(x) > \lambda/2\}, \end{aligned}$$

we have

$$\begin{aligned} & |\{x \in B(0, r) : Mf(x) > \lambda\}| \\ & \leq |\{x \in B(0, r) : Mf_1(x) > \lambda/2\}| + |\{x \in B(0, r) : Mf_2(x) > \lambda/2\}| \\ & \leq C\lambda^{-1} \int_{B(0, 2r)} |f(y)| dy + C|B(0, r)|\lambda^{-1} \int_{\mathbf{R}^n \setminus B(0, 2r)} |f(y)||y|^{-n} dy, \end{aligned}$$

so that

$$\begin{aligned} & r^{-\nu}\lambda|\{x \in B(0, r) : Mf(x) > \lambda\}| \\ & \leq Cr^{-\nu} \int_{B(0, 2r)} |f(y)| dy + Cr^{n-\nu} \int_{\mathbf{R}^n \setminus B(0, 2r)} |f(y)||y|^{-n} dy. \end{aligned}$$

Now it suffices to treat the Hardy type integral in the following:

$$\int_1^\infty \left(r^{n-\nu} \int_{\mathbf{R}^n \setminus B(0, 2r)} |f(y)||y|^{-n} dy \right)^q \frac{dr}{r} < C.$$

In fact, for $\nu < \varepsilon < n$, we note by Hölder's inequality and Fubini's theorem

$$\begin{aligned} & \int_1^\infty \left(r^{n-\nu} \int_{\mathbf{R}^n \setminus B(0, 2r)} |f(y)||y|^{-n} dy \right)^q \frac{dr}{r} \\ & \leq C \sum_j \left(2^{j(n-\nu)} \sum_{k \geq j} 2^{-kn} \int_{B(0, 2^{k+1}) \setminus B(0, 2^k)} |f(y)| dy \right)^q \\ & \leq C \sum_j 2^{j(n-\nu)q} \left(\left(\sum_{k \geq j} 2^{-k(n-\varepsilon)q'} \right)^{q/q'} \sum_{k \geq j} \left(2^{-k\varepsilon} \int_{B(0, 2^{k+1}) \setminus B(0, 2^k)} |f(y)| dy \right)^q \right) \\ & \leq C \sum_j 2^{-j(\nu-\varepsilon)q} \left(\sum_{k \geq j} \left(2^{-k\varepsilon} \int_{B(0, 2^{k+1}) \setminus B(0, 2^k)} |f(y)| dy \right)^q \right) \\ & \leq C \sum_k \left(\left(2^{-k\varepsilon} \int_{B(0, 2^{k+1}) \setminus B(0, 2^k)} |f(y)| dy \right)^q \sum_{j \leq k} 2^{-j(\nu-\varepsilon)q} \right) \\ & \leq C \sum_k \left(2^{-k\nu} \int_{B(0, 2^{k+1}) \setminus B(0, 2^k)} |f(y)| dy \right)^q \\ & \leq C \int_1^\infty \left(r^{-\nu} \int_{B(0, r)} |f(y)| dy \right)^q \frac{dr}{r} \\ & \leq C, \end{aligned}$$

which proves the result. \square

Meskhi [17] obtained the boundedness of the maximal operator in grand Morrey spaces. Here we define grand central Morrey spaces in $L^1(\mathbf{R}^n)$ in the following.

DEFINITION 3.3 (Grand Lebesgue condition). For $1 \leq p \leq \infty$, $0 < q \leq \infty$ and $-\infty < \nu < \infty$, we denote by $M^{p,q,\nu}(\mathbf{R}^n)$ the family of all measurable functions f on \mathbf{R}^n such that

$$\|f\|_{M^{p,q,\nu}(\mathbf{R}^n)} = \sup_{0 < \varepsilon < 1} \left(\int_1^\infty \left(r^{-\nu} \varepsilon \int_{B(0,r)} |f(y)|^{p-\varepsilon} dy \right)^q \frac{dr}{r} \right)^{1/q} < \infty$$

when $q < \infty$ and

$$\|f\|_{M^{p,\infty,\nu}(\mathbf{R}^n)} = \sup_{0 < \varepsilon < 1, r > 1} r^{-\nu} \varepsilon \int_{B(0,r)} |f(y)|^{p-\varepsilon} dy < \infty$$

when $q = \infty$.

It will be expected that the maximal operator M is bounded from $M^{1,q,\nu}(\mathbf{R}^n)$ to $M^{1,q,\nu}(\mathbf{R}^n)$, but the only following weaker result can be proved.

THEOREM 3.4. Let $\nu > 0$ and $0 < q < \infty$. Then there exist constants $A > 0$ and $C > 0$ such that

$$\sup_{0 < \varepsilon < 1} \left(\int_1^\infty \left(r^{-\nu} \varepsilon \int_{\{x \in B(0,r): Mf(x) > 2A\}} \{Mf(x)\}^{1-\varepsilon} dx \right)^q \frac{dr}{r} \right)^{1/q} < C$$

for all f with $\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)} \leq 1$.

4 Generalized Riesz potentials

For $0 < \alpha < n$ and a positive integer k , we define the generalized Riesz potential $I_{\alpha,k}f$ of order α of a locally integrable function f on \mathbf{R}^n by

$$\begin{aligned} I_{\alpha,k}f(x) &= \int_{B(0,1)} I_\alpha(x-y)f(y)dy \\ &\quad + \int_{\mathbf{R}^n \setminus B(0,1)} \left\{ I_\alpha(x-y) - \sum_{\{\lambda: |\lambda| \leq k-1\}} \frac{x^\lambda}{\lambda!} (D^\lambda I_\alpha)(-y) \right\} f(y) dy, \end{aligned}$$

where $I_\alpha(x) = |x|^{\alpha-n}$ (cf. [14, 15]); recall that $I_{\alpha,0}f = I_\alpha f$. Here, $I_\alpha f$ is the Riesz potential of order α of a locally integrable function f on \mathbf{R}^n defined by

$$I_\alpha f(x) = \int_{\mathbf{R}^n} I_\alpha(x-y)f(y)dy.$$

LEMMA 4.1 (cf. [18, 19]). (1) If $2|x| < |y|$, then

$$\left| I_\alpha(x-y) - \sum_{\{\lambda: |\lambda| \leq k-1\}} \frac{x^\lambda}{\lambda!} (D^\lambda I_\alpha)(-y) \right| \leq C|x|^k|y|^{\alpha-n-k}.$$

(2) If $|x|/2 \leq |y| \leq 2|x|$, then

$$\left| I_\alpha(x-y) - \sum_{\{\lambda: |\lambda| \leq k-1\}} \frac{x^\lambda}{\lambda!} (D^\lambda I_\alpha)(-y) \right| \leq C|x-y|^{\alpha-n}.$$

(3) If $1 \leq |y| \leq |x|/2$, then

$$\left| I_\alpha(x-y) - \sum_{\{\lambda: |\lambda| \leq k-1\}} \frac{x^\lambda}{\lambda!} (D^\lambda I_\alpha)(-y) \right| \leq C|x|^{k-1}|y|^{\alpha-n-(k-1)}.$$

Note that $I_{\alpha,k}f$ is finite a.e. on \mathbf{R}^n if

$$\int_{\mathbf{R}^n} (1+|y|)^{\alpha-n-k} |f(y)| dy < \infty.$$

Let p^* denote the Sobolev exponent of p , i.e.,

$$1/p^* = 1/p - \alpha/n.$$

THEOREM 4.2. *Let $\nu \geq 0$, $0 < q \leq \infty$ and k be a positive integer such that $k-1 < \alpha - (n-\nu) < k$. Then the generalized Riesz potential operator $I_{\alpha,k} : f \rightarrow I_{\alpha,k}f$ is bounded from $M^{1,q,\nu}(\mathbf{R}^n)$ to $WM^{1^*,q,\nu}(\mathbf{R}^n)$, that is, there exists a constant $C > 0$ such that*

$$\|I_{\alpha,k}f\|_{WM^{1^*,q,\nu}(\mathbf{R}^n)} \leq C\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)}$$

for $f \in M^{1,q,\nu}(\mathbf{R}^n)$.

Proof. We treat the case $1 < q < \infty$ only.

Let f be a nonnegative measurable function on \mathbf{R}^n such that $\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)} \leq 1$. For $r \geq 1$, we write

$$f = f\chi_{B(0,2r)} + f\chi_{\mathbf{R}^n \setminus B(0,2r)} = f_1 + f_2.$$

If $x \in B(0,r)$, then we have by Lemma 4.1

$$\begin{aligned} |I_{\alpha,k}f_2(x)| &\leq C|x|^k \int_{\mathbf{R}^n \setminus B(0,2r)} |y|^{\alpha-n-k} f(y) dy \\ &\leq Cr^k \int_{\mathbf{R}^n \setminus B(0,2r)} |y|^{\alpha-n-k} f(y) dy \equiv A_1. \end{aligned}$$

On the other hand, for $x \in B(0,r)$,

$$\begin{aligned} |I_{\alpha,k}f_1(x)| &\leq C|x|^{k-1} \int_{B(0,|x|)} |y|^{\alpha-n-(k-1)} f(y) dy + C \int_{B(0,2r)} |x-y|^{\alpha-n} f(y) dy \\ &\leq Cr^{k-1} \int_{B(0,r)} |y|^{\alpha-n-(k-1)} f(y) dy + C \int_{B(0,2r)} |x-y|^{\alpha-n} f(y) dy \\ &\equiv A_2 + CI_\alpha f_1(x), \end{aligned}$$

so that

$$|I_{\alpha,k}f(x)| \leq (A_1 + A_2) + CI_\alpha f_1(x).$$

Now, we set $E = \{x \in B(0,r) : |I_{\alpha,k}f(x)| > \lambda\}$. If $\lambda > 2(A_1 + A_2)$, then

$$\begin{aligned} |E| &\leq |\{x \in E : CI_\alpha f_1(x) > \lambda/2\}| \\ &\leq C\lambda^{-1} \int_E I_\alpha f_1(x) dx \\ &\leq C\lambda^{-1} \int_{\mathbf{R}^n} \left(\int_E I_\alpha(x-y) dx \right) f_1(y) dy \\ &\leq C\lambda^{-1} |E|^{\alpha/n} \int_{\mathbf{R}^n} f_1(y) dy, \end{aligned}$$

so that

$$\lambda|E|^{1/1^*} \leq C \int_{B(0,2r)} f(y)dy,$$

which gives

$$\begin{aligned} & \int_1^\infty (r^{-\nu}\lambda|\{x \in B(0,r) : |I_{\alpha,k}f(x)| > \lambda\}|^{1/1^*})^q \frac{dr}{r} \\ & \leq C \int_1^\infty \left(r^{-\nu} \int_{B(0,2r)} f(y)dy \right)^q \frac{dr}{r} \leq C. \end{aligned}$$

On the other hand, if $\lambda \leq 2(A_1 + A_2)$, then

$$r^{-\nu}\lambda|\{x \in B(0,r) : |I_{\alpha,k}f(x)| > \lambda\}|^{1/1^*} \leq r^{-\nu}2(A_1 + A_2)|B(0,r)|^{1/1^*}.$$

Hence, we treat

$$I_1 = \int_1^\infty \left(r^{-\nu+n/1^*} r^k \int_{\mathbf{R}^n \setminus B(0,2r)} |y|^{\alpha-n-k} f(y)dy \right)^q \frac{dr}{r}$$

and

$$I_2 = \int_1^\infty \left(r^{-\nu+n/1^*} r^{k-1} \int_{B(0,r)} |y|^{\alpha-n-(k-1)} f(y)dy \right)^q \frac{dr}{r}.$$

For I_1 we have

$$\begin{aligned} I_1 & \leq C \sum_i \left(2^{i(-\nu+n-\alpha+k)} \sum_{j \geq i} 2^{j(\alpha-n-k)} \int_{B(0,2^{j+1}) \setminus B(0,2^j)} f(y)dy \right)^q \\ & \leq C \sum_i 2^{i(-\nu+n-\alpha+k)q} \left(\sum_{j \geq i} 2^{-j\varepsilon q'} \right)^{q/q'} \\ & \quad \times \left(\sum_{j \geq i} \left(2^{j(\alpha-n-k+\varepsilon)} \int_{B(0,2^{j+1}) \setminus B(0,2^j)} f(y)dy \right)^q \right) \\ & \leq C \sum_i 2^{i(-\nu+n-\alpha+k-\varepsilon)q} \left(\sum_{j \geq i} \left(2^{j(\alpha-n-k+\varepsilon)} \int_{B(0,2^{j+1}) \setminus B(0,2^j)} f(y)dy \right)^q \right) \\ & \leq C \sum_j \left(\left(2^{j(\alpha-n-k+\varepsilon)} \int_{B(0,2^{j+1}) \setminus B(0,2^j)} f(y)dy \right)^q \left(\sum_{i \leq j} 2^{i(-\nu+n-\alpha+k-\varepsilon)q} \right) \right) \\ & \leq C \sum_j \left(2^{-j\nu} \int_{B(0,2^{j+1}) \setminus B(0,2^j)} f(y)dy \right)^q \leq C \end{aligned}$$

when $0 < \varepsilon < -\nu + n - \alpha + k$. Similarly, for I_2 we have

$$I_2 \leq C \sum_j \left(2^{-j\nu} \int_{B(0,2^{j+1}) \setminus B(0,2^j)} f(y)dy \right)^q \leq C$$

when $0 < \varepsilon < \nu - n + \alpha - k + 1$. Thus

$$\int_1^\infty (r^{-\nu}\lambda|\{x \in B(0,r) : |I_{\alpha,k}f(x)| > \lambda\}|^{1/1^*})^q \frac{dr}{r} \leq C,$$

which completes the proof. \square

REMARK 4.3. Suppose $p > 1$ and $1/p - \alpha/n > 0$ and $k - 1 < \alpha - (n - \nu)/p < k$. In view of [22, Theorem 4.5], one can find a constant $C > 0$ such that

$$\|I_{\alpha,k}f\|_{M^{p^*,\infty,\nu}(\mathbf{R}^n)} \leq C\|f\|_{M^{p,\infty,\nu}(\mathbf{R}^n)}$$

for all $f \in M^{p,\infty,\nu}(\mathbf{R}^n)$. The case $0 < q < \infty$ can be treated in a way similar to the above proof.

REMARK 4.4. Let $\nu \geq 0$ and $0 < q < \infty$. Suppose $\alpha - (n - \nu)$ is a nonnegative integer $k - 1$. Then there exists a constant $C > 0$ such that

$$\sup_{\lambda>0, r>1} \left(\int_1^\infty (r^{-\nu}(\log(1+r))^{-1}\lambda|\{x \in B(0,r) : |I_{\alpha,k}f(x)| > \lambda\}|^{1/1^*})^q \frac{dr}{r} \right)^{1/q} < C$$

for all f with $\|f\|_{M^{1,q,\nu}(\mathbf{R}^n)} \leq 1$.

References

- [1] D. R. Adams and J. Xiao, Morrey spaces in harmonic analysis, *Ark. Mat.* **50** (2012), no. 2, 201–230.
- [2] A. Almeida and D. Drihem, Maximal, potential and singular type operators on Herz spaces with variable exponents, *J. Math. Anal. Appl.* **394** (2012), no. 2, 781–795.
- [3] A. Almeida, J. Hasanov and S. Samko, Maximal and potential operators in variable exponent Morrey spaces, *Georgian Math. J.* **15** (2008), 195–208.
- [4] J. Alvarez, M. Guzmán-Partida and J. Lakey, Spaces of bounded λ -central mean oscillation, Morrey spaces, and λ -central Carleson measures, *Collect. Math.* **51** (2000), 1–47.
- [5] A. Beurling, Construction and analysis of some convolution algebras, *Ann. Inst. Fourier* **14** (1964), 1–32.
- [6] V. I. Burenkov, A. Gogatishvili, V. S. Guliyev and R. Ch. Mustafayev, Boundedness of the fractional maximal operator in local Morrey-type spaces, *Complex Var. Elliptic Equ.* **55** (2010), no. 8-10, 739–758.
- [7] V. I. Burenkov, A. Gogatishvili, V. S. Guliyev and R. Ch. Mustafayev, Boundedness of the Riesz potential in local Morrey-type spaces, *Potential Anal.* **35** (2011), no. 1, 67–87.
- [8] V. I. Burenkov and H. V. Guliyev, Necessary and sufficient conditions for boundedness of the maximal operator in the local Morrey-type spaces, *Stud. Math.* **163**(2) (2004), 157–176.
- [9] V. I. Burenkov, H. V. Guliyev and V. S. Guliyev, On boundedness of the fractional maximal operator from complementary Morrey-type spaces to Morrey-type spaces. The interaction of analysis and geometry, 1732, *Contemp. Math.*, 424, Amer. Math. Soc., Providence, RI, 2007.

- [10] A. Fiorenza and G. E. Karadzhov, Grand and small Lebesgue spaces and their analogs. *Z. Anal. Anwendungen* **23** (2004), no. 4, 657–681.
- [11] Z. Fu, Y. Lin and S. Lu, λ -central BMO estimates for commutators of singular integral operators with rough kernels, *Acta Math. Sin. (Engl. Ser.)* **24** (2008), no. 3, 373–386.
- [12] V. Kokilashvili, A. Meskhi, and H. Rafeiro, Riesz type potential operators in generalized grand Morrey spaces, *Georgian Math. J.* **20** (2013), 43–64.
- [13] C. Herz, Lipschitz spaces and Bernstein’s theorem on absolutely convergent Fourier transforms, *J. Math. Mech.* **18** (1968), 283–324.
- [14] K. Matsuoka, Generalized fractional integrals on central Morrey spaces and generalized λ -CMO spaces, in *Function Spaces X*, Banach Center Publ., 102, 181188, Inst. Math., Polish Acad. Sci., Warszawa, 2014.
- [15] K. Matsuoka, Generalized fractional integrals on central Morrey spaces and generalized σ -Lipschitz spaces, in *Current Trends in Analysis and its Applications: Proceedings of the 9th ISAAC Congress, Kraków 2013*, 179189, Birkhäuser Basel, 2015.
- [16] K. Matsuoka and E. Nakai, Fractional integral operators on $B^{p,\lambda}$ with Morrey-Campanato norms, *Function Spaces IX* (Krakow, Poland, 2009), 249–264, Banach Center Publications, Vol.92, Institute of Mathematics, Polish Academy of Sciences, Warszawa, 2011.
- [17] A. Meskhi, A, Maximal functions, potentials and singular integrals in grand Morrey spaces, *Complex Var. Elliptic Equ.* **56**:10–11 (2011), 1003–1019.
- [18] Y. Mizuta, *Potential theory in Euclidean spaces*, Gakkōtoshō, Tokyo, 1996.
- [19] Y. Mizuta, Integral representations, differentiability properties and limits at infinity for Beppo Levi functions, *Potential Analysis* **6** (1997), 237–276.
- [20] Y. Mizuta and T. Ohno, Sobolev’s theorem and duality for Herz-Morrey spaces of variable exponent, *Ann. Acad. Sci. Fenn. Math.* **39** (2014), 389–416.
- [21] Y. Mizuta and T. Ohno, Herz-Morrey spaces of variable exponent, Riesz potential operator and duality, *Complex Var. Elliptic Equ.* **60** (2015), no. 2, 211–240.
- [22] Y. Mizuta, T. Ohno and T. Shimomura, Boundedness of maximal operators and Sobolev’s theorem for non-homogeneous central Morrey spaces of variable exponent, *Hokkaido Math. J.* **44** (2015), 185–201.
- [23] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, 1970.

College of Economics

Nihon University

1-3-2 Misaki-cho Kanda Chiyoda-ku Tokyo 101-8360, Japan

E-mail : katsu.m@nihon-u.ac.jp

and

4-13-11 Hachi-Hom-Matsu-Minami

Higashi-Hiroshima 739-0144, Japan

E-mail : yomizuta@hiroshima-u.ac.jp